

Efficacy of magnetic stimulation for female stress urinary incontinence: a meta-analysis

Kai Sun^{*} , Dongxu Zhang^{*}, Gang Wu, Tianqi Wang, JiTao Wu, Hongxu Ren and Yuanshan Cui

Ther Adv Urol

2021, Vol. 13: 1–12

DOI: 10.1177/
17562872211032485

© The Author(s), 2021.
Article reuse guidelines:
sagepub.com/journals-
permissions

Abstract

Aim: This meta-analysis aimed to evaluate the efficacy of magnetic stimulation (MS) in treating female stress urinary incontinence (SUI) and providing an alternative treatment for patients who are unwilling to undergo surgery.

Methods: Randomized controlled trials (RCTs) that evaluated MS as a remedy for female SUI were retrieved from various electronic databases, including MEDLINE, EMBASE, and the Cochrane Controlled Trial Registry system. Moreover, reference lists for related papers were carefully screened for relevant studies.

Results: A total of six RCTs evaluating the effect of MS in treating female SUI were included in this study. Compared with the placebo group, the MS group exhibited higher quality-of-life scores [mean difference (MD) 0.59, 95% credibility interval (CI) 0.23–0.95; $p=0.001$] and lower International Consultation on Incontinence Questionnaire scores (MD –3.93, 95% CI –5.85 to –2.01; $p<0.0001$). Moreover, they exhibited a higher objective cure rate (odds ratio 8.49, 95% CI 3.08–23.37). In addition, MS treatment reduced the number of episodes of urinary incontinence (MD –1.42, 95% CI –2.24 to –0.59; $p=0.0007$) and urine loss on pad test (MD –4.67, 95% CI –8.05 to –1.28; $p=0.007$). There were no significant treatment-related adverse reactions.

Conclusion: This study evaluated the efficacy and safety of MS in the treatment of female SUI. The results have important implications for patients who do not wish to undergo surgical therapy. We found that MS treatment for SUI has positive outcomes, however, future studies should aim at establishing the best protocol for optimizing the therapeutic effect.

Keywords: magnetic stimulation, meta-analysis, randomized controlled trials, stress urinary incontinence

Received: 6 April 2021; revised manuscript accepted: 25 June 2021.

Introduction

Urinary incontinence (UI), defined by the International Continence Society and International Urogynecological Association as involuntary urination, is a common, chronic, and distressing disorder that lessens the quality of life (QoL), much like chronic diseases such as stroke.^{1–3} Among the three main types of UI identified by the Standardization Steering Committee, the most common is stress urinary incontinence (SUI).⁴ Even though its incidence rate varies among different locations, it has exhibited a tremendous

yearly increase, which causes significant negative economic and social impacts.^{5,6}

Conservative and surgical therapies are the main options for managing female SUI. Burch and urethral sling procedures are regarded as primary surgical procedures with a high cure rate of between 70% and 90%.⁷ Due to complications associated with invasive procedures, such as pelvic pain and difficulties in urinating, however, surgery is not a preferable therapeutic option. According to the 2017 European Association of Urology guidelines

Correspondence to:
Yuanshan Cui
Urology Department,
The Affiliated Yantai
Yuhuangding Hospital of
Qingdao University, No. 20
East Yuhuangding Road,
Yantai, Shandong 264000,
China
s1009471740@163.com

Hongxu Ren
Operation Department,
The Affiliated Yantai
Yuhuangding Hospital of
Qingdao University, No. 20
East Yuhuangding Road,
Yantai, Shandong 264000,
China
s979429197@163.com

Kai Sun
Dongxu Zhang
Gang Wu
Tianqi Wang
JiTao Wu
Urology Department,
The Affiliated Yantai
Yuhuangding Hospital of
Qingdao University, Yantai,
Shandong, China

^{*}These authors contributed equally to this work.

on UI therapy, pelvic floor muscle training (PFMT), bladder training, electrical stimulation (ES), magnetic stimulation (MS), and posterior tibial nerve stimulation are potential therapeutic options for SUI.⁸ PFMT, which has been proven to be effective in previous randomized controlled trials (RCTs),^{9–12} has been recommended as the initial therapeutic option for SUI by the American Urological Association (AUA) and Society of Urodynamics, Female Pelvic Medicine & Urogenital Reconstruction (SUFU). Improvement in clinical outcomes for SUI after PFMT range from 50% to 70%; however, due to poor compliance, the cure rate does not exceed 15–30%.^{13,14} Notably, ES, a replacement therapy, has a success rate of 48–70%.¹⁵ Due to the discomfort or pain associated with high-intensity percutaneous currents, the use of ES as a therapeutic option is limited.^{16–20} In 1998, the US Food and Drug Administration recommended MS as the first choice therapeutic option for SUI.²¹ Since then, clinical applications for MS have improved due to its associated security, automatic contraction, absence of malaise from probe insertion, and ease of administration.

Various clinical trials have evaluated the efficacy of MS in ameliorating female SUI with positive outcomes. One systematic review²² evaluated the effect of MS on the treatment of SUI, but did not adequately provide the number of included RCTs, analysis of outcome indicators, or the therapeutic mechanism of MS. The above analyses only involved studies written in English, which potentially could have omitted several other relevant studies. We, therefore, conducted a meta-analysis, which not only summarized the data from published reviews, but also integrated the data from other newly published RCTs to evaluate comprehensively the effects of MS on SUI in terms of improving QoL and reducing the incidence and severity of UI. By summarizing reports in other relevant literature, we further discuss the mechanism of MS in the treatment of SUI and put forward reasonable suggestions for the treatment plan.

Materials and methods

Search strategy

This meta-analysis was performed in accordance with the Preferred Reporting Items for Systematic

Reviews and Meta-Analyses (PRISMA).²³ All RCTs evaluating the association between SUI and MS, and published up to 1 March 2021 were independently reviewed by two authors. Appropriate trials were extracted from electronic databases, including MEDLINE, EMBASE, and the Cochrane Controlled Trials Register, using various combinations of Medical Subject Heading terms. The search terms used were ‘magnetic stimulation’, ‘stress urinary incontinence’, and ‘RCTs’. Repeated studies were excluded from analysis. A third party was involved to mediate any arising disputes. There were no language restrictions for the included studies.

Inclusion criteria and trial selection

The inclusion criteria for the RCTs were: (a) those evaluating the curative effect of MS as a remedy for female SUI; (b) those with full content and relevant data that could be acquired; (c) those with authentic data, chiefly incorporating the sum of subjects and the meritorious consequences of each index. Provided the same outcomes were issued in various journals or at diverse times, the updated research results were absorbed in this meta-analysis. Results from the same group of researchers were included once.

Quality assessment

The Jadad score and the Cochrane bias risk assessment tool were used to evaluate the methodological quality of all included RCTs.^{24,25} Generation of randomization sequences and incomplete outcome data were involved in quality standardization. Moreover, blinding, allocation concealment, along with freedom from selective reporting and other biases, were incorporated (Table 1).

Data extraction

Two reviewers used predefined data extraction forms to independently extract data. Disagreements were resolved through mediation by a senior author. The extracted data included: (a) year of publication, first author’s name and country of origin; (b) the type of remedy to which participants had access; (c) therapeutic plan; (d) SUI results; (e) duration of follow up; (f) the total number of participants in each group.

Table 1. Details of included studies.

Study, country	Sample size (n) MS/sham	Age, years (median)	Inclusion criteria	Exclusion criteria	MS	Intervention			Length of intervention period	Follow-up period	Outcome measures	Quality assessment
						Location	Intensity	Frequency				
Fujishiro <i>et al.</i> , ²⁶ Japan	31/31	58	≥ 1 episode of leakage recorded in a voiding diary; ≥ 2g urine loss in 1-h pad test	Disorders causing any LUTS; ongoing treatment for SUI	Sacral roots (S3)	50% of maximum output	15 Hz 5 S/min	30 min	Once only	1 week	1. Maximum urethral closure pressure and cytometry; 2. No. of leakages; 3. Pad test/g [24h]; 4. QoL scores	High risk
Manganotti <i>et al.</i> , ²⁷ Italy	10/10	50.1	≥ 1 leakage recorded in a 3-day voiding diary; ≥ 2g urine loss in a 1-h pad test or a positive standardized stress test	Disorders causing LUTS; severe cardiac or cerebrovascular disorders; receiving treatment for SUI	Sacral roots (S2–S4)	60% of maximum output	15 Hz 3 S/min	15 min	Three sessions per week for 2 weeks	1 month	1. QoL scores; 2. Pad test/g [24h]; 3. Standardized stress test	Low risk
Gilling <i>et al.</i> , ²⁸ New Zealand	35/35	54.4	Symptoms of SUI; genuine SUI by pad-testing and urodynamics; neurologically normal; stable detrusor function on urodynamics with cystometric capacity of >200 ml, PFR >10 ml/s, PVR <100 ml	Previous incontinence or pelvic floor surgery; grade 3 or 4 POP; pregnant; drugs for bladder dysfunction; internal devices with electrical or magnetic component; pelvic or lower limb metallic prosthesis	Pelvic floor	Maximum level tolerated by the patient	10 Hz; 50 Hz	A 10-min stimulation at 10 Hz; a 3-min rest; a 10-min stimulation at 50 Hz.	Three treatment sessions per week for 6 weeks	6 months	1. 20-min pad test; 2. 3-day bladder diary; 3. 24-h pad test; 4. No. of pads used; 5. PFM strength	Low risk
Tsai <i>et al.</i> , ²⁹ Taipei China	14/10	63.1	A diagnosis of SUI, with or without detrusor overactivity, confirmed by urodynamic results; a SUI history of at least 6 months, which remained refractory after at least 1 month of first-line management	History of surgery or hormone replacement therapy for SUI; severe pelvic prolapse (>grade 3 prolapse or Qmax <15 ml/s); contraindications for SMS; received anticholinergic medication	Sacral roots (S3)	Maximum level tolerated by the patient	5 Hz; in 10-s on/20-s off cycles	20 min	12 consecutive week days	18 weeks	1. Cystometric; 2. UPP; 3. U-UDI; 4. OAB-Q	Low risk

(continued)

Table 1. (continued)

Study, country	Sample size (n) MS/sham	Age, years (median)	Inclusion criteria	Exclusion criteria	MS	Location			Length of intervention period	Follow-up period	Outcome measures	Quality assessment
						Intensity	Frequency	Duration				
Lim <i>et al.</i> ³⁰ , Malaysia	60/60	52.5	Female aged 21 or older with urine leak upon coughing; a ICIQ-UI SF score of 6 points or greater; can perform the 1-h pad test	Other subtypes of UI; pelvic irradiation; contraindications for MS; previous surgery for SUJ; previous treatment with PMS; prolapse stage III or IV; severe urethral sphincter weakness or urethral/vesical fistula; post-void residual volume greater than 200 ml; pregnancy	Pelvic floor	Maximum level tolerated by the patient	50 Hz; in 8-s on/4-s off cycles	20 min	Two sessions per week for 2 months	14 months	1. ICIQ scores; 2. Pad test/g (1 h); 3. No. of leakages; 4. 1-h pad test; 5. PFM function; 6. PGI-I; 7. ICIQ-LUTS QoL	Low risk
Yamanishi ³² , Japan	18/12	NA	Women with urodynamic SUJ refractory to PFMT for more than 12 weeks and who did not want to undergo surgery	UI due to detrusor overactivity; complications after pelvic surgery or trauma; wearing a pacemaker; complicated by malignancy; with a residual urine volume \geq 200 ml	Pelvic floor	Maximum level tolerated by the patient	50 Hz; in 5-s on/5-s off cycles	20 min	One session per week for 10 weeks	10 weeks	1. No. of leakages; 2. Pad test/g (24 h); 3. QoL scores; 4. ICIQ scores; 5. ALPP	Low risk

ALPP, abdominal leak point pressure; ICIQ, International Consultation on Incontinence Questionnaire; ICIQ-LUTS QoL, ICIQ-lower urinary tract symptoms quality of life; ICIQ-UI SF, ICIQ-urinary incontinence-short form; LUTS, lower urinary tract symptoms; MS, magnetic stimulation; NA, not available; PFMT, pelvic floor muscle training; OAB-Q, overactive bladder questionnaire; PFM, pelvic floor muscle; PFR, peak flow rate; PGI-I, patient global impression of improvement; PMS, pulsed magnetic stimulation; POP, pelvic organ prolapse; PVR, post-void residual; QoL, quality of life; SMS, simultaneous multislice imaging; SUJ, stress urinary incontinence; UI, urinary incontinence; UPP, urethral pressure profile; U-UDI, urge-urinary distress inventory.

Statistical analysis

Review Manager version 5.3.0 (Cochrane Collaboration, Oxford, UK) was used for data analysis. Fixed- or random-effect models were adopted for appraising indicators. Mean difference (MD) was used to interpret continuous data and odds ratios (ORs) for dichotomous outcomes, coupled with 95% CI.³¹ The I-square (I^2) test was used to evaluate the effect of heterogeneity on the meta-analysis results. In cases where the I^2 value was greater than 50%, a random-effects model was used, whereas if the I^2 value was less than 50%, a fixed-effects model was used to evaluate the data. $p \leq 0.05$ was set as the threshold for statistical significance.

Results

Characteristics of the trials

Based on the inclusion and exclusion criteria, a total of 352 articles were extracted. However, after a review of all titles and abstracts, a total of 89 articles were excluded. Due to lack of data, a total of 25 articles were excluded from the remaining 32 articles. Two reviewers separately rated the absolute papers and made a selection following the criteria. Of the remaining 25 articles, 14 articles were excluded for not using appropriate outcome indicators, 6 articles were excluded for not being RCTs, and 5 articles were excluded for using subgroups that were not suitable for analysis. Finally, seven articles containing six RCTs assessing the efficacy of MS for female patients with SUI^{26–30,32,33} were included in this study. The selection and elimination PRISMA flowchart is presented in Figure 1. Characteristics of the studies are presented in Table 1. The risk of bias graph and summary are shown in Figure 2.

QoL scores

QoL scores were extracted from 6 articles involving 336 participants (174 in the MS group and 162 in the sham group). We used a random-effects model to evaluate these RCTs, but the I^2 test implied heterogeneity (Figure 3(a)). The study by Lim *et al.*³³ had the greatest effect. Eliminating this study and using the fixed-effects model remarkably reduced the I^2 to 39%. The MD was 0.59, while the 95% CI was 0.23–0.95 ($p=0.001$) (Figure 3(b)), implying that MS therapy improved QoL. Then, we performed subgroup analysis of the location of MS to understand the impact on QoL scores, which showed an MD of




Figure 1. Flowchart of the study selection process. RCT, randomized controlled trial.

0.59, whereas the 95% CI was 0.00–1.18 ($p=0.05$) in sacral roots and the MD was 2.7 while the 95% CI was 0.15–5.25 ($p=0.04$) in the pelvic floor (Figure 3(c)). These findings indicate that the effect was slightly different due to different stimulation positions of the MS and sham groups.




Figure 2. (a) Risk of bias summary: review authors' judgements about each risk of bias item for each included study. (b) Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

Pad test

A total of 5 RCTs involving 197 participants (107 in the MS group and 90 in the sham group) had their pad-test data. Statistical heterogeneity was high, I^2 of 47% ($p=0.42$) (Figure 4(a)). Given the high heterogeneity between studies, we performed sensitivity analysis. The cross-sectional trial of Manganotti *et al.*²⁷ was the only one that was removed. After omission of the study, we found an MD of -4.67 and 95% CI of -8.05 to -1.28 ($p=0.007$), without heterogeneity (Figure 4(b)). Patients in the MS treatment group had significantly less urine loss on pad test than those in the placebo group.

Leaks

Three studies involving 127 participants (72 in the MS group and 55 in the sham group)

presented their data on the sum of leaks/week through a voiding diary. A fixed-effects model was selected for analysis. Compared with the sham group, the MS group exhibited a valid decline (MD -1.42 ; 95% CI -2.24 to -0.59 ; $p=0.0007$), and there was no heterogeneity (Figure 4(c)).

ICIQ scores

Pooled RCTs involving 185 participants (101 in the MS group and 84 in the sham group) presented data on ICIQ scores. A fixed-effects model was introduced to rate these RCTs, notably, the MD was -3.93 and 95% CI was -5.85 to -2.01 ($p<0.0001$) (Figure 4(d)). These findings show that ICIQ scores were lower in MS-treated patients.

Objective cure rate

Pooled RCTs with data for objective cure (leakage less than 1 g on the 1-h pad test) rate were used to evaluate the improvement in incontinence symptoms. Patients treated with MS were more likely to be associated with a higher objective cure rate (OR 8.49, 95% CI 3.08 23.37; $p<0.0001$) (Figure 4(e)).

Discussion

Despite the small sample sizes and non-uniform treatment regimens used in the majority of the tests, pooled analysis of the data showed that in terms of overall efficacy, the efficacy of MS for SUI was always superior to that of the sham group.

QoL scores, as the most popular indicator for evaluating female SUI treatment, was greatly improved in the MS group compared with the sham group. Studies that assessed QoL reported encouraging results. In their study, Hoşcan *et al.*³⁴ reported that the mean QoL score increased from 61.6 to 75.4 after MS ($p=0.003$). Elsewhere, Lo *et al.*³⁵ used the Urge-Urinary Distress Inventory (U-UDI) to measure QoL. They found an improvement in total UDI-6 scores. The King's Health Questionnaire is also popular when calculating QoL.³⁶ Irrespective of which questionnaire was used to evaluate QoL, results tended to be positive. In addition, the ICIQ score questionnaire is highly recommended by the 5th ICI.³⁷ Due to the rise in in-depth and precise




Figure 3. Forest plot comparing the change in (a) QoL scores, (b) QoL scores after omitting study, (c) QoL scores in subgroup analysis of the location of MS between active and sham groups.

CI, confidence interval; df, degrees of freedom; MS, magnetic stimulation; SD, standard deviation.

experiments, more meaningful data will emerge, which will enhance our judgement on the efficacy of MS.

Even though studies by Gilling *et al.*²⁸ and Yamanishi *et al.*³² did not report effective findings, whereas that by Manganotti *et al.*²⁷ showed high heterogeneity, pooled data exhibited a positive result in urine loss on pad test. In a single report, deteriorative outcomes in 35.5% of women with SUI based on pad tests were found.³⁸

However, the study lacked motivation and had a high dropout rate of 35.4%. Another study concluded that the 24-h pad test had no advantage in

predicting diacrisis of SUI.³⁹ A potential breakthrough however was reported by Hoşcan *et al.*³⁴ who found that pad weight was reduced from 14.4 ± 10.7 g to 6.5 ± 5.1 g at 3 months in the MS group. Notably, the pad test has many detection schemes, which may lead to deviations in measurement results. Generally, the pad test is controversial and is not a stable test indicator, therefore, more studies are needed to establish the best detection process.

The frequency of incontinence is a vital objective indicator for assessing the reliability of MS treatment. Although our meta-analysis included three




Figure 4. (a) Forest plot comparing the change in (a) pad test, (b) pad test after the omitting study, (c) number of leaks, (d) ICIQ scores, (e) objective cure rate between the active and sham groups. CI, confidence interval; df, degrees of freedom; ICIQ, International Consultation on Incontinence Questionnaire; MS, magnetic stimulation; SD, standard deviation.

RCTs, pooled data showed that the reduction in frequency of SUI was statistically significant. Galloway *et al.*⁴⁰ found a significant reduction in the median number of pads, as well as a significant reduction in leakage events and frequency of detrusor instability, which concurs with our

findings and supports the effectiveness of MS in reducing the frequency of incontinence.

A large proportion of patients with moderate or below SUI mostly present with external urethral sphincter (EUS) and pelvic floor muscle

weakness.⁴¹ EUS has complete neuromuscular innervation and can be used to cope with movement by increasing its size and strength.⁴² Eddy currents can be induced by transcutaneous MS in the pelvis from where they flow into tissues, thereby depolarizing the axons. As a peripheral motor nerve axon, the impulse passes to the motor endplate, inducing the mandatory release of acetylcholine. Then, homologous muscle fibers depolarize and contract.^{40,43} MS may modify the activity in pelvic floor muscle groups, as well as the discharge pattern and frequency of motor nerve fibers responsible for resting tension of the pelvic floor and sphincter. Moreover, MS is associated with a significant increase in bladder volume, which may be attributed to acute activation of the inhibitory detrusor reflex pathway after stimulation of the pudendal afferent nerve. Fujishiro *et al.*²⁶ and Tsai *et al.*²⁹ reported changes in bladder volume and maximum urethral closure pressure after MS treatment. Bladder capacity in the MS group was significantly higher than that in the sham operation group. Determination of maximum urethral closing pressure, however, did not draw the same conclusion. Elsewhere, Fujishiro *et al.*²⁶ reported that the maximum urethral closure pressure did not increase, however, their findings were disapproved by Tsai *et al.*²⁹ who concluded that it did. This may be correlated to the different methods of measurement used in the two studies.

The low level of standardization of the MS protocol is challenging. Different studies used different stimulus intensities, frequencies, locations, and durations. Until now, the optimal frequency and duration of the pulse remain controversial. It has been shown that frequencies of 20–50 Hz are effective for SUI, and satisfactory pelvic floor contraction during SUI treatment requires a higher dose of 50 Hz.^{44,45} Therefore, the treatment may not be as effective as expected in three RCTs using stimuli at frequencies ranging from 5 Hz to 15 Hz. We performed subgroup analysis of QoL scores for the two different stimulation sites of sacral roots and pelvic floor, and the results revealed little difference between the two groups. Based on the above findings, we preliminarily suggested that the stimulation site might affect the therapeutic effect, and that the pelvic floor may exhibit better outcomes. However, studies have not evaluated the effects of different stimulus sites. Therefore, there is a need to determine

whether outcome indicators can be improved by stimulating other parts, apart from sacral roots and pelvic floor. Treatment and follow-up duration varied across studies, which inevitably led to differences in outcomes. Galloway *et al.*⁴⁶ concluded that active MS of the pelvic floor twice a week for 6 weeks improved SUI, which remained effective after 3 months. It has also been reported that the benefits of MS worsen over time,^{40,46,47} perhaps because of the treatment regimen. Therefore, to improve the effectiveness of MS treatment and to promote MS in clinical treatment, an appropriate MS protocol must be developed.

In addition, patients subjected to MS therapy were found to exhibit a higher objective cure rate. It has been reported that the cure rate and improvement rate after Electromagnetic stimulation (EMS) treatment were not much better than those after PFMT treatment.⁴² Moreover, Hoşcan *et al.*³⁴ reported a 29.7% cure rate. However, after 3 months, they reported a 48.1% improvement rate, as well as an extraordinary improvement in QoL. Suzuki *et al.*⁴⁸ reported a cure rate of 20% in the MS group after active treatment. The results could not be used to disparage MS treatment for patients who were all non-responders to PFMT or drug therapy. This can be explained by active contractions produced by the PFMT exercise muscle strength, which is better than passive contraction induced by MS. Therefore, the therapeutic effect is not satisfactory.

Reported side effects of MS, including lower limbs, abdominal, and back pain, among others, were not severe or life threatening.³⁸ Fewer side effects associated with MS were reported in the included articles, suggesting that it is relatively safe and tolerable.

This meta-analysis has some limitations. We noted that the quality of the included studies was flawed, in terms of study designs, patient selection, blinding, publication bias, and outcome data. Our results are based on unadjusted estimates. More accurate results will originate from adjustments of other confounders, such as gender, body mass index, lifestyle, and age among others. More RCTs with abundant sample sizes are needed to validate our findings. Additional RCTs should be performed to ascertain the virtue and defects of MS in treating female SUI.

Conclusion

We found that MS may be beneficial in the clinical management of female SUI, especially for patients who do not consent to surgery. More clinical trials are needed to determine the appropriate protocol for optimizing the therapeutic effect.

Author contributions

KS and GW: designing the study; analyzing the extracted data; writing the first manuscript draft.

DZ: searching the literature; extracting and analyzing the data.

TQW and JW: analyzing the data; providing critical scientific input.

HR and YC: resolving discrepancies about the quality of the included studies; reviewing the manuscript; providing critical scientific input.

All authors approved the final version of this manuscript.

Conflict of interest statement

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was supported by grants from the National Natural Science Foundation of China (Nos. 81870525; 81572835), Taishan Scholars Program of Shandong Province (No. tsqn201909199).

ORCID iD

Kai Sun  <https://orcid.org/0000-0001-8866-8838>

References

1. Bo K, Frawley HC, Haylen BT, *et al.* An International Urogynecological Association (IUGA)/International Continence Society (ICS) joint report on the terminology for the conservative and nonpharmacological management of female pelvic floor dysfunction. *Neurourol Urodyn* 2017; 36: 221–244.
2. Horng SS, Huang N, Wu SI, *et al.* The epidemiology of urinary incontinence and its influence on quality of life in Taiwanese middle-aged women. *Neurourol Urodyn* 2013; 32: 371–376.
3. Shulz SE and Kopec JA. Impact of chronic conditions. *Health Rep* 2003; 14: 41.
4. National Collaborating Centre for Women's and Children's Health. *Urinary incontinence in women: the management of urinary incontinence in women.* London: Royal College of Obstetricians and Gynaecologists, 2013, pp.906–911.
5. Irwin DE, Mungapen L, Milsom I, *et al.* The economic impact of overactive bladder syndrome in six Western countries. *BJU Int* 2009; 103: 202–209.
6. Subak LL, Goode PS, Brubaker L, *et al.* Urinary incontinence management costs are reduced following Burch or sling surgery for stress incontinence. *Am J Obstet Gynecol* 2014; 211: 171 e171–e177.
7. Bergman A, Koonings PP and Ballard CA. Primary stress urinary incontinence and pelvic relaxation: prospective randomized comparison of three different operations. *Am J Obstet Gynecol* 1989; 161: 97–101.
8. Lucas MG, Bosch RJ, Burkhard FC, *et al.* [European Association of Urology guidelines on assessment and nonsurgical management of urinary incontinence.] *Actas Urol Esp* 2013; 37: 199–213.
9. Bo K, Talseth T and Holme I. Single blind, randomised controlled trial of pelvic floor exercises, electrical stimulation, vaginal cones, and no treatment in management of genuine stress incontinence in women. *BMJ* 1999; 318: 487–493.
10. Felicissimo MF, Carneiro MM, Saleme CS, *et al.* Intensive supervised versus unsupervised pelvic floor muscle training for the treatment of stress urinary incontinence: a randomized comparative trial. *Int Urogynecol J* 2010; 21: 835–840.
11. Gormley EA, Lightner DJ, Faraday M, *et al.*; American Urological A, Society of Urodynamics FPM. Diagnosis and treatment of overactive bladder (non-neurogenic) in adults: AUA/SUFU guideline amendment. *J Urol* 2015; 193: 1572–1580.
12. Takahashi S, Takei M, Nishizawa O, *et al.* Clinical guideline for female lower urinary tract symptoms. *Low Urin Tract Symptoms* 2016; 8: 5–29.
13. Elia G and Bergman A. Pelvic muscle exercises: when do they work? *Obstet Gynecol* 1993; 81: 283–286.
14. Klarskov P, Belving D, Bischoff N, *et al.* Pelvic floor exercise versus surgery for female urinary stress incontinence. *Urol Int* 1986; 41: 129–132.

15. Miller K, Richardson DA, Siegel SW, *et al.* Pelvic floor electrical stimulation for genuine stress incontinence: who will benefit and when? *Int Urogynecol J Pelvic Floor Dysfunct* 1998; 9: 265–270.
16. Yokoyama T, Nishiguchi J, Watanabe T, *et al.* Comparative study of effects of extracorporeal magnetic innervation versus electrical stimulation for urinary incontinence after radical prostatectomy. *Urology* 2004; 63: 264–267.
17. Correia GN, Pereira VS, Hirakawa HS, *et al.* Effects of surface and intravaginal electrical stimulation in the treatment of women with stress urinary incontinence: randomized controlled trial. *Eur J Obstet Gynecol Reprod Biol* 2014; 173: 113–118.
18. Wang S, Lv J, Feng X, *et al.* Efficacy of electrical pudendal nerve stimulation versus transvaginal electrical stimulation in treating female idiopathic urgency urinary incontinence. *J Urol* 2017; 197: 1496–1501.
19. Yamanishi T, Kaga K, Fuse M, *et al.* Neuromodulation for the treatment of lower urinary tract symptoms. *Low Urin Tract Symptoms* 2015; 7: 121–132.
20. Yamanishi T, Kamai T and Yoshida K. Neuromodulation for the treatment of urinary incontinence. *Int J Urol* 2008; 15: 665–672.
21. Goldberg RP and Sand PK. Electromagnetic pelvic floor stimulation for urinary incontinence and bladder disease. *Int Urogynecol J Pelvic Floor Dysfunct* 2001; 12: 401–404.
22. Lim R, Lee SW, Tan PY, *et al.* Efficacy of electromagnetic therapy for urinary incontinence: a systematic review. *Neurourol Urodyn* 2015; 34: 713–722.
23. Moher D, Liberati A, Tetzlaff J, *et al.* Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 2009; 62: 1006–1012.
24. Jadad AR, Moore RA, Carroll D, *et al.* Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996; 17: 1–12.
25. Cumpston M, Li T, Page MJ, *et al.* Updated guidance for trusted systematic reviews: a new edition of the Cochrane Handbook for Systematic Reviews of Interventions. *Cochrane Database Syst Rev* 2019; 10: ED000142.
26. Fujishiro T, Enomoto H, Ugawa Y, *et al.* Magnetic stimulation of the sacral roots for the treatment of stress incontinence: an investigational study and placebo-controlled trial. *J Urol* 2000; 164: 1277–1279.
27. Manganotti P, Zaina F, Vedovi E, *et al.* Repetitive magnetic stimulation of the sacral roots for the treatment of stress incontinence: a brief report. *Eura Medicophys* 2007; 43: 339–344.
28. Gilling PJ, Wilson LC, Westenberg AM, *et al.* A double-blind randomized controlled trial of electromagnetic stimulation of the pelvic floor vs sham therapy in the treatment of women with stress urinary incontinence. *BJU Int* 2009; 103: 1386–1390.
29. Tsai PY, Wang CP, Hsieh CY, *et al.* Long-term sacral magnetic stimulation for refractory stress urinary incontinence. *Arch Phys Med Rehabil* 2014; 95: 2231–2238.
30. Lim R, Liong ML, Leong WS, *et al.* Pulsed magnetic stimulation for stress urinary incontinence: 1-year followup results. *J Urol* 2017; 197: 1302–1308.
31. DerSimonian R and Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; 7: 177–188.
32. Yamanishi T, Suzuki T, Sato R, *et al.* Effects of magnetic stimulation on urodynamic stress incontinence refractory to pelvic floor muscle training in a randomized sham-controlled study. *Low Urin Tract Symptoms* 2019; 11: 61–65.
33. Lim R, Liong ML, Leong WS, *et al.* Effect of pulsed magnetic stimulation on quality of life of female patients with stress urinary incontinence: an IDEAL-D stage 2b study. *Int Urogynecol J* 2018; 29: 547–554.
34. Hoşcan MB, Dilmen C, Perk H, *et al.* Extracorporeal magnetic innervation for the treatment of stress urinary incontinence: results of two-year follow-up. *Urol Int* 2008; 81: 167–172.
35. Lo TS, Tseng LH, Lin YH, *et al.* Effect of extracorporeal magnetic energy stimulation on bothersome lower urinary tract symptoms and quality of life in female patients with stress urinary incontinence and overactive bladder. *J Obstet Gynaecol Res* 2013; 39: 1526–1532.
36. Kelleher CJ, Cardozo LD, Khullar V, *et al.* A new questionnaire to assess the quality of life of urinary incontinent women. *Br J Obstet Gynaecol* 1997; 104: 1374–1379.
37. Avery K, Donovan J, Peters TJ, *et al.* ICIQ: a brief and robust measure for evaluating the symptoms and impact of urinary incontinence. *Neurourol Urodyn* 2004; 23: 322–330.
38. Ismail SI, Forward G, Bastin L, *et al.* Extracorporeal magnetic energy stimulation of pelvic floor muscles for urodynamic stress

- incontinence of urine in women. *J Obstet Gynaecol* 2009; 29: 35–39.
39. Henderson JW, Kane SM, Mangel JM, *et al.* A randomized comparative study evaluating various cough stress tests and 24-hour pad test with urodynamics in the diagnosis of stress urinary incontinence. *J Urol* 2018; 199: 1557–1564.
40. Galloway NT, El-Galley RE, Sand PK, *et al.* Update on extracorporeal magnetic innervation (EXMI) therapy for stress urinary incontinence. *Urology* 2000; 56(6 Suppl. 1): 82–86.
41. Hunskaar S, Arnold EP, Burgio K, *et al.* Epidemiology and natural history of urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct* 2000; 11: 301–319.
42. Fowler SB. Pelvic floor muscle training versus no treatment, or inactive control treatments, for urinary incontinence in women. *Clin Nurse Spec* 2011; 25: 226–227.
43. Quek P. A critical review on magnetic stimulation: what is its role in the management of pelvic floor disorders? *Curr Opin Urol* 2005; 15: 231–235.
44. Janez J, Plevnik S and Suhel P. Urethral and bladder responses to anal electrical stimulation. *J Urol* 1979; 122: 192–194.
45. Yamanishi T, Yasuda K, Suda S, *et al.* Effect of functional continuous magnetic stimulation for urinary incontinence. *J Urol* 2000; 163: 456–459.
46. Galloway NT, El-Galley RE, Sand PK, *et al.* Extracorporeal magnetic innervation therapy for stress urinary incontinence. *Urology* 1999; 53: 1108–1111.
47. Yokoyama T, Fujita O, Nishiguchi J, *et al.* Extracorporeal magnetic innervation treatment for urinary incontinence. *Int J Urol* 2004; 11: 602–606.
48. Suzuki T, Yasuda K, Yamanishi T, *et al.* Randomized, double-blind, sham-controlled evaluation of the effect of functional continuous magnetic stimulation in patients with urgency incontinence. *Neurourol Urodyn* 2007; 26: 767–772.